International Journal of Modern Physics A
© World Scientific Publishing Company

arXiv:0907.1128v1 [hep-th] 7 Jul 2009

STATIC WORMHOLES IN VACUUM FOR CONFORMAL GRAVITY

JULIO OLIVA

Centro de Estudios Científicos (CECS), Casilla 1469, Valdivia, Chile.
Instituto de Física, Universidad Austral de Chile, Casilla 567, Valdivia, Chile.

DAVID TEMPO

Centro de Estudios Científicos (CECS), Casilla 1469, Valdivia, Chile.
Departamento de Física, Universidad de Concepción, Casilla 160-C, Concepción, Chile.
Physique théorique et mathématique, Université Libre de Bruxelles, ULB Campus Plaine C.P.231, B-1050 Bruxelles, Belgium.

RICARDO TRONCOSO

Centro de Estudios Científicos (CECS), Casilla 1469, Valdivia, Chile.
Centro de Ingeniería de la Innovación del CECS (CIN), Valdivia, Chile.

A static spherically symmetric wormhole solution for conformal gravity in vacuum is found. The solution possesses a single integration constant which determines the size of the neck connecting two static homogeneous universes of constant spatial curvature. Time runs at different rates on each side of the neck, and depending on the value of the parameter, the wormhole can develop a cosmological horizon only at one side. It is shown that the wormholes correspond to the matching of different Einstein spacetimes by means of improper conformal transformations.

Keywords: Wormholes; Conformal gravity; Charge without charge.

PACS numbers: 04.20.Jb, 04.50.Kd

If one wonders about the possibility of realizing static wormholes in vacuum in four dimensions, one knows that it should be performed within some scenario beyond General Relativity (see e.g. Ref. 1). Experience gathered from extensions of General Relativity in diverse dimensions[2–10] suggests that one of the key features allowing this possibility is the relaxation of the asymptotic conditions for the metric. An interesting theory possessing this last feature is conformal gravity (see e.g. Refs. 11,12,13), whose Lagrangian can be written as the square of the Weyl Tensor, so that the action reads

$$I = \alpha \int d^4x\sqrt{-g}\; C^{\alpha\beta\mu\nu}C_{\alpha\beta\mu\nu}\ . \tag{1}$$

Apart from being invariant under diffeomorphisms, this action is also invariant under local rescalings of the metric given by $g_{\mu\nu} \to \Omega^2(x)g_{\mu\nu}$, and the field equations

are given by the vanishing of the Bach tensor, i.e.,

$$\left(\nabla^{\mu}\nabla^{\nu}+\frac{1}{2}R^{\mu\nu}\right)C^{\alpha}{}_{\mu\beta\nu}=0\ . \tag{2}$$

It is simple to verify that any Einstein space with cosmological constant is a solution of the theory in vacuum.

Conformal gravity was intensively studied in the past and it has been shown to be renormalizable[14] – Note that the coupling α is dimensionless. Nevertheless, the theory possesses fourth-order field equations for the metric, so that it has ghosts as it is generically expected. In the context we are interested in, regardless the theory is suitable or not as the ultimate one to describe gravity, it certainly deserves to be studied.

It is also worth pointing out that it has been recently shown that for smooth matter distributions, wormholes that do not violate the weak energy condition near the throat can exist in conformal gravity.[15] For our purposes, it is useful mentioning that the most general spherically symmetric solution of conformal gravity (see e.g. Ref. 16) possesses a relaxed asymptotic behavior as compared with General Relativity. Thus, for the reasons explained above, the door is open to look for wormholes in vacuum within this theory.

It can be seen that conformal gravity admits the following static spherically symmetric wormhole solution in vacuum, whose metric reads

$$ds^2=-\left(1+a^2\tanh\rho\right)dt^2+\frac{d\rho^2}{1+a^2\tanh\rho}+l_0^2\cosh^2\rho\ d\Omega^2\ , \tag{3}$$

where $d\Omega^2$ stands for the line element of S^2, and the range of the radial coordinate is given by $-\infty<\rho<\infty$. The wormhole possesses a single integration constant a that parameterizes the radius of the neck given by $l_0^2=\left(3a^4+1\right)^{-1/2}$, being located at $\rho=0$. The wormhole connects two static homogeneous universes of constant spatial curvature with different radii, as it can be seen from the asymptotic behavior of the curvature

$$R^{mn}{}_{kl}\underset{\rho\to\pm\infty}{\longrightarrow}-\left(1\pm a^2\right)\delta^{mn}{}_{kl}\ , \tag{4}$$

and time runs at different rates at each side of the neck, since

$$g_{tt}\underset{\rho\to\pm\infty}{\longrightarrow}-\left(1\pm a^2\right)\ . \tag{5}$$

The case of $a^2\to-a^2$ is obtained from reflexion symmetry of the radial coordinate.

For $a^2<1$ the wormhole interpolates between static universes with spatial geometries given by hyperbolic spaces of radii $\left(1\pm a^2\right)^{-1/2}$. In the case of $a=0$ the metric acquires a simple form, given by

$$ds^2=-dt^2+d\rho^2+\cosh^2\rho\ d\Omega^2\ , \tag{6}$$

which at both sides is asymptotically locally described by $R \times H_3$, and it can be regarded as "the groundstate". For $a = 1$ the wormhole interpolates between flat space and a static universe with spatial geometry given by a hyperbolic space of radius $2^{-1/2}$. In the case of $a^2 > 1$ the wormhole develops a cosmological horizon at one side of the neck, located at $\rho = \rho_c = -\tanh^{-1}\left(1/a^2\right)$, and interpolates between an Einstein Universe ($R \times S^3$) for $\rho \to -\infty$, and $R \times H_3$.

The causal structure of the wormhole coincides with the one of Minkowski spacetime in two dimensions for $a^2 \leq 1$. For $a^2 > 1$ the wormhole shares the same causal structure with two-dimensional Rindler spacetime, as shown in Fig. 1a.
It can also be shown that the wormholes in vacuum generically correspond to the matching of different Einstein spacetimes at infinity by means of improper conformal transformations. A particularly interesting case corresponds to $a^2 > 1$ since the wormhole is conformally related to the matching of the patch covering the region within the event horizon and infinity of the Schwarzschild-de Sitter (SdS) metric with positive mass with two additional patches obtained from the region beyond the cosmological horizon of the Schwarzschild-de Sitter metric with negative mass. This is depicted in Fig. 1b. Note that in General Relativity the matching of different spaces through the boundary of their corresponding conformal compactifications has no sense, since the proper distance to pass from one patch to the other diverges. Nonetheless, it is worth pointing out that within the context of conformal gravity this kind of matching can be performed by means of an improper conformal transformation, i.e., a conformal transformation that vanishes at the matching surface. In this way, the proper distance required to pass through a pair of points located at each side of the matching surface, given by $ds^2 = \Omega^2 ds_E^2$, becomes finite.

It can also be seen that the wormhole (3) is conformally related by patches to the spherically symmetric solution of conformal gravity found in Refs. 17 and 16. This feature also holds for the generalization of the wormholes presented here, having necks of genus greater than one. The extension of the wormhole (3) with radial electric or magnetic fields can also be found, and it turns out to have electric or magnetic "charge without charge". For further details we refer to Ref. 7. As an ending remark, it is worth mentioning that the definition of mass in conformal gravity is a very subtle issue, which is not free of controversy.[18–19] Thus, in order to have a suitable analysis of the solutions, the construction of finite conserved charges written as suitable surface integrals for conformal gravity should be addressed.

Acknowledgments

We thank the organizers of the "7th Alexander Friedmann International Seminar on Gravitation and Cosmology" for the nice atmosphere provided at the meeting. Special thanks to Gastón Giribet, Carlos Kozameh, Claude LeBrun and Roberto Troncoso for very useful comments. This work was partially funded by FONDECYT grants 1095098, 1061291, 1071125, 1085322, 3085043; David Tempo thanks CONICYT and Escuela de Graduados of the Universidad de Concepción for finan-

cial support. The Centro de Estudios Científicos (CECS) is funded by the Chilean Government through the Millennium Science Initiative and the Centers of Excellence Base Financing Program of CONICYT. CECS is also supported by a group of private companies which at present includes Antofagasta Minerals, Arauco, Empresas CMPC, Indura, Naviera Ultragas and Telefónica del Sur. CIN is funded by CONICYT and the Gobierno Regional de Los Ríos.

References

1. M. Visser, *Lorentzian wormholes: From Einstein to Hawking*, Woodbury, USA: AIP (1995).
2. G. Dotti, J. Oliva and R. Troncoso, *Phys. Rev. D* **75**, 024002 (2007).
3. E. Gravanis and S. Willison, *Phys. Rev. D* **75**, 084025 (2007).
4. G. Dotti, J. Oliva and R. Troncoso, *Phys. Rev. D* **76**, 064038 (2007).
5. G. Dotti, J. Oliva and R. Troncoso, *Preprint CECS-PHY-07/21.*
6. C. Garraffo, G. Giribet, E. Gravanis and S. Willison, *J. Math. Phys.* **49**, 042502 (2008).
7. J. Oliva, D. Tempo and R. Troncoso, *Preprint CECS-PHY-08/09.*
8. H. Maeda and M. Nozawa, *Phys. Rev. D* ***78,*** 024005 (2008).
9. D. H. Correa, J. Oliva and R. Troncoso, *J. High Energy Phys.* **0808**, 081 (2008).
10. F. Canfora and A. Giacomini, Phys. Rev. D **78**, 084034 (2008).
11. E. S. Fradkin and A. A. Tseytlin, *Phys. Rept.* **119**, 233 (1985).
12. N. Boulanger and M. Henneaux, *Annalen Phys.* **10**, 935 (2001).
13. N. Berkovits and E. Witten, *J. High Energy Phys.* **0408**, 009 (2004).
14. K. S. Stelle, *Phys. Rev. D***16**, 953 (1977); *Gen. Rel. Grav.* **9**, 353 (1978).
15. F. S. N. Lobo, *Class. Quant. Grav.* ***25***, 175006 (2008).
16. R. J. Riegert, *Phys. Rev. Lett.* **53**, 315 (1984).
17. R. Bach *Math. Z.* **9**, 110 (1921).
18. D. G. Boulware, G. T. Horowitz, and A. Strominger, *Phys. Rev. Lett.* 50, 1726 (1983).
19. S. Deser and B. Tekin, *Phys. Rev. Lett.* **89**, 101101 (2002); *Phys. Rev. D* **67**, 084009 (2003); *Phys. Rev. D* **75**, 084032 (2007).



Fig. 1. (a) Corresponds to the causal structure of the wormhole with $a > 1$. (b) Shows how this wormhole is constructed from the conformal matching of two SdS spacetimes with positive and negative masses.